



ASCLERA (polidocanol) Injection is the **ONLY FDA-approved** liquid polidocanol 0.5% and 1% formulations used for the treatment of uncomplicated spider and reticular veins in the lower legs.<sup>1</sup>

# ASCLERA®

(Polidocanol) Injection

**FDA approved to treat spider and reticular leg veins.**

ASCLERA is contraindicated for patients with known allergy to polidocanol and patients with acute thromboembolic diseases. In clinical studies, the following adverse reactions were observed after using ASCLERA and were more common with ASCLERA than placebo: injection site hematoma, injection site irritation, injection site discoloration, injection site pain, injection site itching, injection site warmth, neovascularization, injection site clotting.

Please consult the package insert for full prescribing information including warnings and precautions, available on request by calling Methapharm Medical Information at 1-866-701-4636, or to view and download at [www.ASCLERA.com](http://www.ASCLERA.com).

To report SUSPECTED ADVERSE REACTIONS, contact Methapharm Medical Information at 1-866-701-4636, or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

For all other ASCLERA inquiries please call 1-833-766-8346.




Visit [www.ASCLERA.com](http://www.ASCLERA.com) for more information THEN ask your Healthcare Provider about ASCLERA to begin your sclerotherapy treatment today.

**methapharm**  
Specialty Pharmaceuticals

Distributed by Methapharm, Inc. Copyright © 2025 Methapharm, Inc. All rights reserved. ASCLERA® is a registered trademark of Chemische Fabrik Kreussler & Co. GmbH.

ASC-US-VPBR1-1225



Asc for  
**ASCLERA®**  
(Polidocanol) Injection

# ASCLERA®

(Polidocanol) Injection

## Simple procedure for sustained results

Uncomplicated spider vein treatment ( $\leq 1$  mm)  
patient treated with 0.5% ASCLERA



Before



After

Uncomplicated reticular vein treatment (1-3 mm)  
patient treated with 1% ASCLERA



Before



After

Improvement 18 weeks  
after last treatment

the **Gold Standard** in sclerotherapy of small veins

## What to expect before and after your treatment

### DURING<sup>1</sup>

- ⌚ Treatment typically lasts between 15-45 minutes.
- ✨ Antiseptic is used to clean the area.
- 💉 A very fine needle is used to inject ASCLERA.
- 🌀 A slight sting may accompany each injection.

NOTE: depending on the severity of the veins treated, you may require additional treatments.

### AFTER<sup>1</sup>

- 🚶 Immediately following treatment, you will need to walk for 15-20 minutes.
- 🧦 You will need to wear compression stockings for:
  - **Spider veins:** 2 to 3 days after treatment
  - **Reticular veins:** 5 to 7 days after treatment

NOTE: Speak to your physician about length of time to wear compression stockings during the daytime (possible 2-3 weeks during the day).

### Avoid for 2-3 days after treatment:

- 🏋 Heavy exercise
- 🧖 Hot tubs and saunas
- ✈ Long flights
- ☀ Sunbathing

84%

of patients treated with ASCLERA were satisfied after 26 weeks compared to 63% treated with a competitive product.<sup>2</sup>

87%

of patients were satisfied or very satisfied with their ASCLERA treatment at 12 weeks<sup>2,†</sup>

95%

of patients treated with ASCLERA showed good or complete treatment success as rated by physicians.<sup>2</sup>

**Ask your Healthcare Provider about ASCLERA to begin your sclerotherapy treatment today. Revitalizing and maintaining your legs begins NOW.**

### References:

- 1) ASCLERA® Full Prescribing Information. Methapharm, Inc.
- 2) Rabe E, Schliephake D, Otto J, Breu FX, et al. Sclerotherapy of telangiectases and reticular veins: a double-blind, randomized, comparative clinical trial of polidocanol, sodium tetradecyl sulphate and isotonic saline (EASI study). *Phlebology* 2010;25:124-31.